




# The Need for Hadronic Data in the Resonance Region




Michael Sadler  
Abilene Christian University  
Abilene, Texas USA



# Nucleon model states and $N\pi$ couplings

BW Mass  
(MeV/c<sup>2</sup>)



# N(1440) summary in 2004 Review of Particle Physics

Citation: S. Eidelman et al. (Particle Data Group), Phys. Lett. B 592, 1 (2004) (URL: <http://pdg.lbl.gov>)

**N(1440) P<sub>11</sub>**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

## N(1440) BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1430 to 1470 (≅ 1440) OUR ESTIMATE</b>			
1462±10	MANLEY	92	IPWA πN → πN & Nππ
1440±30	CUTKOSKY	80	IPWA πN → πN
1410±12	HOEHLER	79	IPWA πN → πN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1518±5	PENNER	02c	DPWA Multichannel
1479±80	VRANA	00	DPWA Multichannel
1463±7	ARNDT	96	IPWA γN → πN
1467	ARNDT	95	DPWA πN → Nπ
1421±18	BATINIC	95	DPWA πN → Nπ, Nη
1465	LI	93	IPWA γN → πN
1471	CUTKOSKY	90	IPWA πN → πN
1411	CRAWFORD	80	DPWA γN → πN
1472	<sup>1</sup> BAKER	79	DPWA π <sup>-</sup> p → nη
1417	BARBOUR	78	DPWA γN → πN
1460	BERENDS	77	IPWA γN → πN
1380	<sup>2</sup> LONGACRE	77	IPWA πN → Nππ
-----	-----	-----	-----

# N(1440) summary in 2008 Review of Particle Physics

Citation: C. Amsler *et al.* (Particle Data Group), PL **B667**, 1 (2008) (URL: <http://pdg.lbl.gov>)

**$N(1440) P_{11}$**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

Most of the results published before 1975 were last included in our 1982 edition, Physics Letters **111B** 1 (1982). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006).

## N(1440) BREIT-WIGNER MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

**1420 to 1470 ( $\approx 1440$ ) OUR ESTIMATE**

1485.0 $\pm$ 1.2	ARNDT	06	DPWA $\pi N \rightarrow \pi N, \eta N$
1462 $\pm$ 10	MANLEY	92	IPWA $\pi N \rightarrow \pi N \& N\pi\pi$
1440 $\pm$ 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1410 $\pm$ 12	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1436 $\pm$ 15	SARANTSEV	08	DPWA Multichannel
1468.0 $\pm$ 4.5	ARNDT	04	DPWA $\pi N \rightarrow \pi N, \eta N$
1518 $\pm$ 5	PENNER	02C	DPWA Multichannel
1479 $\pm$ 80	VRANA	00	DPWA Multichannel

# N\*(1710) summary in Review of Particle Physics

Citation: W.-M. Yao et al. (Particle Data Group), J. Phys. G **33**, 1 (2006) and 2007 partial update for edition 2008 (URL: <http://pdg.lbl.gov>)

**N(1710) P<sub>11</sub>**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 were last included in our 1982 edition, *Physics Letters* **111B** 1 (1982). Some further obsolete results published before 1980 were last included in our 2006 edition, *Journal of Physics*, G **33** 1 (2006).

The latest GWU analysis (ARNDT 06) finds no evidence for this resonance.

## N(1710) BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1680 to 1740 (as 1710) OUR ESTIMATE</b>			
1717 ± 28	MANLEY	92	IPWA πN → πN & Nππ
1700 ± 50	CUTKOSKY	80	IPWA πN → πN
1723 ± 9	HOEHLER	79	IPWA πN → πN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1752 ± 3	PENNER	02c	DPWA Multichannel
1699 ± 65	VRANA	00	DPWA Multichannel
1720 ± 10	ARNDT	96	IPWA γN → πN
1766 ± 34	<sup>1</sup> BATINIC	95	DPWA πN → Nπ, Nη
1706	CUTKOSKY	90	IPWA πN → πN
1692	CRAWFORD	80	DPWA γN → πN
1730	SAXON	80	DPWA π <sup>-</sup> p → ΛK <sup>0</sup>
1720	<sup>2</sup> LONGACRE	77	IPWA πN → Nππ
1710	<sup>3</sup> LONGACRE	75	IPWA πN → Nππ

# N\*(1710) summary in Review of Particle Physics

## N(1710) DECAY MODES


The following branching fractions are our estimates, not fits or averages.

	Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$	$N\pi$	10–20 %
$\Gamma_2$	$N\eta$	( 6.2 ± 1.0 ) %
$\Gamma_3$	$N\omega$	(13.0 ± 2.0) %
$\Gamma_4$	$\Lambda K$	5–25 %
$\Gamma_5$	$\Sigma K$	
$\Gamma_6$	$N\pi\pi$	40–90 %
$\Gamma_7$	$\Delta\pi$	15–40 %
$\Gamma_8$	$\Delta(1232)\pi, P\text{-wave}$	
$\Gamma_9$	$N\rho$	5–25 %
$\Gamma_{10}$	$N\rho, S=1/2, P\text{-wave}$	
$\Gamma_{11}$	$N\rho, S=3/2, P\text{-wave}$	
$\Gamma_{12}$	$N(\pi\pi)_{S\text{-wave}}^{J=0}$	10–40 %
$\Gamma_{13}$	$p\gamma$	0.002–0.05%
$\Gamma_{14}$	$p\gamma, \text{ helicity}=1/2$	0.002–0.05%
$\Gamma_{15}$	$n\gamma$	0.0–0.02%
$\Gamma_{16}$	$n\gamma, \text{ helicity}=1/2$	0.0–0.02%

# Dynamical origin of P11 resonances

Suzuki, Julia-Diaz, Kamano, Lee, Matsuyama, Sato, PRL104 042302 (2010)

All three P11 poles below 2 GeV are generated from a *same, single* bare state!



*Multi-channel* reactions can generate *many* resonance poles from a *single* bare state  
Eden, Taylor, Phys. Rev. 133 B1575 (1964)


Evidences in hadron and nuclear physics are summarized e.g., in Morgan, Pennington PRL59 2818 (1987)



(This slide courtesy of Igor Strakovsky)

# $N^*$ and $\Delta^*$ States Coupled to $\pi N$

- One of the most convincing ways to study Spectroscopy of  $N^*$  &  $\Delta^*$  is  $\pi N$  PWA
- Main objects in the PDG Listings [<http://pdg.lbl.gov/>] come from: Karlsruhe-Helsinki, Carnegie-Mellon-Berkeley, and GW/VPI
- GW DAC SAID program:  $\pi N \rightarrow \pi N \Rightarrow \gamma N \rightarrow \pi N \Rightarrow \gamma^* N \rightarrow \pi N$ 
  - $\pi N$  elastic amplitudes from fits to the observables:  $\sigma^{\text{tot}}$ ,  $d\sigma/d\Omega$ , and  $P$  plus a few  $R$  and  $A$  measurements, 0.4 %
  - Contains resonances contributing to  $\gamma^* N \rightarrow \pi N$
  - Assuming dominance of 2 hadronic channels, can parametrize  $\gamma^* N \rightarrow \pi N$  in terms of  $\pi N \rightarrow \pi N$  amplitudes alone
  - Resulting multipoles can be re-fitted in terms of Res/Bckgr contributions or used as input to multi-channel fits with more elaborate constraints
  - Comparison of various resonance-extraction methods gives a more reliable estimate of systematic (model) errors



(This slide courtesy of Igor Strakovsky)

# Summary of $N^*$ and $\Delta^*$ from GWU08 Analysis







[R. Arndt, W. Briscoe, I. Strakovsky, R. Workman, Phys Rev C **74**, 045205 (2006)]

- Standard PWA reveals only **wide** Resonances, but not too wide ( $\Gamma < 500$  MeV) and possessing **not too small** BR (BR  $> 4\%$ )
- Standard PWA (by construction) tends to miss **narrow** Resonances with  $\Gamma < 30$  MeV
- Our study **does not** support several  $N^*$  and  $\Delta^*$  reported by PDG2008:
  - \*\*\*  $\Delta(1600)P_{33}$ ,  $N(1700)D_{13}$ ,  $N(1710)P_{11}$ ,  $\Delta(1920)P_{33}$
  - \*\*  $N(1900)P_{13}$ ,  $\Delta(1900)S_{31}$ ,  $N(1990)F_{17}$ ,  $\Delta(2000)F_{35}$ ,  $N(2080)D_{13}$ ,  
 $N(2200)D_{15}$ ,  $\Delta(2300)H_{39}$ ,  $\Delta(2750)I_{313}$
  - \*  $\Delta(1750)P_{31}$ ,  $\Delta(1940)D_{33}$ ,  $N(2090)S_{11}$ ,  $N(2100)P_{11}$ ,  $\Delta(2150)S_{31}$ ,  
 $\Delta(2200)G_{37}$ ,  $\Delta(2350)D_{35}$ ,  $\Delta(2390)F_{37}$
- Our study **does** suggest several 'new'  $N^*$  and  $\Delta^*$ :
  - \*\*\*\*  $\Delta(2420)H_{311}$
  - \*\*\*  $\Delta(1930)D_{35}$ ,  $N(2600)I_{111}$  [BW, no Pole]
  - \*\*  $N(2000)F_{15}$ ,  $\Delta(2400)G_{39}$
  - new  $N(2245)H_{111}$  [CLAS ?]



# Summary of Pion-Nucleon Reactions and $\pi N$ Experiments at LAMPF and BNL

## Reactions




## Observables

$$d\sigma/d\Omega$$

$$P (=A_N)$$

A and R

## Comparison of the spin rotation parameter A with PWA predictions



# Summary of Present PWA Efforts

$\pi N \rightarrow \gamma N \rightarrow$	$\pi N$	$\pi\pi N$	$\eta N$	$K\Lambda, K\Sigma$
SAID	● ●	●	● ●	
Bonn-Gatchina	● ●	● ●	● ●	● ●
EBAC	● ●	● ●	● ★	★ ★
Juelich-UGA	● ●	● ★	●	● ★
Zagreb	● ★	● ★	● ★	● ★
Giessen	● ●	● ●	● ●	● ●

★ In progress

Adapted from T. Sato review talk, "EBAC meeting", Jefferson Laboratory, May 2010,  
and B. Julia-Diaz, MENU2010, College of William and Mary, June 2010.



# Summary of Present PWA Status

- Listings in Review of Particle Properties rely primarily on Karlsruhe-Helsinki and Carnegie Mellon-Berkeley PWAs from 1980.
- GWU (formerly VPI) PWA includes recent data, but differs significantly in prediction of resonances.
- New efforts by EBAC, Bonn-Gatchina, Juelich, Zagreb and Giessen are producing results but are ignored in RPP.
- To date, no “missing resonances” found! Indeed, have we ‘lost’ the  $P_{11}(1710)$ ?
- Analyses should include  $\pi p \rightarrow \pi\pi N$ , new data are needed.
- **Hadronic data are needed to analyze data from photo- and electro-production of resonances.**


# $\pi^- p \rightarrow \pi^- p$ and $\pi^- p \rightarrow K\Lambda$ at ITEP

An Existing Facility for Measurements in Baryon Spectroscopy  
Institute for Theoretical and Experimental  
Physics, Moscow, Russia

I.G. Alekseev, P.Ye. Budkovsky, V.P. Kanavets, M.M. Kats, L.I. Koroleva,  
V.V. Kulikov, B.V. Morozov, V.M. Nesterov, V.V. Ryltsov, V.A. Sakharov,  
A.D. Sulimov, D.N. Svirida, **ITEP**

A.I. Kovalev, N.G. Kozlenko, V.S. Kozlov, A.G. Krivshich, D.V. Novinsky, V.V.  
Sumachev, V.Yu. Trautman, Ye.A. Filimonov, **PNPI**

M.E. Sadler, J. Kish, D. Soboyede, E. Walker, S. Watson **ACU**






# Project at ITEP (EPECUR)

- Measurements of  $\pi^-p \rightarrow \pi^-p$  and  $\pi^-p \rightarrow K\Lambda$  at  $P_\pi = 900 - 1200 \text{ MeV}/c$  ( $\sqrt{s} = 1610 - 1770 \text{ MeV}$ )
- Participating Institutions are ITEP, PNPI and ACU
- Emphasis on narrow resonance search and  $N^*$  (1710) (second  $P_{11}$ )
- Natural extension of LAMPF, PNPI and BNL (Crystal Ball) programs
- Preparation for experiments at J-PARC or FAIR?




# $\pi^-p \rightarrow \pi^-p$ at ITEP

- Differential cross sections at 40-120° CM as function of the invariant mass of  $\pi^-p$ -system.
- “Formation”-type experiment: invariant mass resolution (0.7 MeV) is based on the high momentum resolution (0.1%) of the magneto-optic channel.
- Statistical resolution as high as 0.5 %
- Obtain clear evidence for a narrow (2-20 MeV) resonance even if its elasticity is only 1%.
- Main parts of experimental setup are liquid hydrogen target and proportional and drift chambers.




# Sensitivity for $\pi^- p \rightarrow \pi^- p$

- Momentum range 900-1200 MeV/c, 40-120° CM
- $\sqrt{s} = 1610-1770$  MeV
- Invariant mass intervals of 0.5 MeV
- Statistical precision of 0.5%
- Started data taking in 2009



$\pi^- p, P_{\text{lab}} = 1060 \text{ MeV/c}$


С учетом эффективностей и акцептанса



# Strangeness Production ( $\Lambda^0$ )


- $\pi^- p \rightarrow K^0 \Lambda$ 
  - ↳  $\pi^- p$
  - ↳  $K_S \rightarrow \pi^+ \pi^-$
- Important reaction because:
  - Sizable cross section.
  - Pure  $I = 1/2$  selects only  $N^*$  resonances.
- Little is known about resonances that decay to  $K \Lambda$ .
- Will also be able to determine the final-state  $\Lambda$  polarization, since it is self-analyzing (two observables).
- Precise differential cross section and polarization data for  $\pi^- p \rightarrow K^0 \Lambda$  should be straightforward to analyze with a PWA.

# Current situation – experimental data for $\pi^-p \rightarrow K^0\Lambda$




# $\pi^- p \rightarrow K \Lambda$ at ITEP


- Differential cross-section with statistical precision 1% and step in the invariant mass 0.5 MeV at the angles 0-180° CM.
- Momentum range 900-1200 MeV/c  $\Rightarrow$  1610-1770 MeV
- ~24 days of running, after  $\pi^- p \rightarrow \pi^- p$  measurements




# GEANT4 Simulation for $\pi^-p \rightarrow K\Lambda$



# Resonance formation and decay for $\pi^- p \rightarrow K^0 \Lambda$








# New Possibility for Measurements in Baryon Spectroscopy

## The Main Injector Particle Production Experiment (MIPP-FNAL-P960) at Fermilab



## *MIPP Upgrade collaboration list*

*M. Sadler*

Abilene Christian University

*V. Singh*

Banaras Hindu University, Varanasi 221005, India

*H. Kirk, R.B. Palmer,*

Brookhaven National Laboratory, Upton, New York

*M.R. Anantharaman, V.C. Kuriakose, M. Sabir, R.B. Thayyullathil*

Cochin University of Science and technology, Kochi 682022, India

*R.J. Peterson*

University of Colorado, Boulder

*B.C. Choudhary*

University of Delhi, Delhi 110007, India

*W. Baker, B. Baldin, D. Carey, D. Christian, M. Demarteau, D. Jensen, C. Johnstone,  
R. Raja, A. Ronzhin, N. Solomey, W. Wester, J-Y Wu, R. Zwaska*

Fermi National Accelerator Laboratory

*Bill Briscoe, Igor Strakovsky, Ron Workman*

George Washington University, Washington D.C

*H. Gutbrod, B. Kolb, K. Peters,*

GSI, Darmstadt, Germany

*G. Feldman,*

Harvard University

*B. Bambah, R. Mohanta, E. Harikumar*

University of Hyderabad, India

*A.K. Giri,*

Indian Institute of Technology, Hyderabad, Andhra Pradesh 502205, India

*Y. Torun,*

Illinois Institute of Technology

*M. Messier, J. Paley*

Indiana University

*U. Akgun, G. Aydin, F. Duru, E. Gülmez, Y. Gunaydin, Y. Onel, A. Penzo*

University of Iowa



## *MIPP Upgrade collaboration list*

V.Avdeichicov,R.Leitner,J.Manjavidze,V.Nikitin,I.Rufanov,T.Topuria  
Joint Institute for Nuclear Research, Dubna, Russia

D.M.Manley,

Kent State University

H.Löhner, J.Messchendorp,  
KVI, Groningen, Netherlands

H.R.Gustafson,M.Longo,T.Nigmanov, D.Rajaram

University of Michigan

V.Bhatnagar, A.Kumar, S. Mahajan, S. Sahijpal, A. Singh  
Panjab University, Chandigarh, 160014, India

S.P.Kruglov,I.V.Lopatin,N.G.Kozlenko,A.A.Kulbardis,D.V.Nowinsky, A.K.Radkov,V.V.Sumachev  
Petersburg Nuclear Physics Institute, Gatchina, Russia

K. McDonald,

Princeton University, Princeton, New Jersey

A.Bujak, L.Gutay,

Purdue University

D.Bergman, G.Thomson

Rutgers University

A.Godley,S.R.Mishra,C.Rosenfeld

University of South Carolina

C.Dukes,C.Materniak,K.Nelson,A.Norman

University of Virginia

P.Desiati, F.Halzen, T.Montaruli,

University of Wisconsin, Madison

P.Sokolsky, W.Springer

University of Utah

H.Meyer, N.Solomey,

Wichita State University, Kansas


18 new institutions have joined. 6 from India.

New Collaborators in color cyan

# MIPP

## Main Injector Particle Production Experiment (FNAL-E907)

Horizontal cut plane



## TPC installation





# MIPP-TPC

- The Time Projection Chamber was built by the BEVALAC group at LBL for heavy ion studies in 1990's. Donated to Fermilab after usage at BNL. It took approximately \$3 million to construct.
- Can handle high multiplicity events. Time to drift across TPC=16  $\mu$ s.
- Electronic equivalent of bubble chamber, high acceptance, with dE/dx capabilities. Dead time 16ms. Unreacted beam swept out in 8ms. Can tolerate  $10^5$  particles per second going through it.
- TPC dimensions 96 x 75 x 150 cm.
- Previous data taking rate was ~60 Hz. Electronics upgraded to increase rate to ~1000 Hz.



# TPC processed event

MPP (FNAL E907)



Run: 9303  
 SubRun: 0  
 Event: 1

Thu Aug 19 2004  
 02:52:42.951283

Version: 0  
 Trigger: 10008709



h31













## Extension of MIPP


- Pion beams as low as 1 GeV/c possible with new magnet controls.
- Was proposed to Fermilab PAC in 2006. Approval was deferred until data analysis of first run is completed and running in collider mode is ended. Was rejected in November 2010.
- If eventually approved, opens possibility of making measurements in the  $\pi N$  resonance region.
- ACU (M. E. Sadler), Kent State University (M. Manley), and PNPI (V. V. Sumachev, S. P. Kruglov, I. V. Lopatin, N. G. Kozlenko, A. A. Kulbardis, and D. V. Nowinsky) have agreed to participate in new proposal.
- Additional collaborators are invited!








# 657 MeV/c $\pi^- p \rightarrow \pi^0 n$



(This slide courtesy of Igor Strakovsky)  
 Electromagnetic Probe for Resonance Physics





## Single pion production



- Single pion production is an important reaction to search for missing resonances that couple weakly to  $\pi N$ .
- With a TPC, one can measure 4 of the 5 reactions amenable to  $\pi^\pm p$  scattering:  
 $\pi^+ p \rightarrow \pi^+ \pi^0 p$  and  $\pi^- p \rightarrow \pi^- \pi^0 p$  (detect  $\pi^0$  by missing mass)  
 $\pi^+ p \rightarrow \pi^+ \pi^+ n$  and  $\pi^- p \rightarrow \pi^- \pi^+ n$  (detect  $n$  by missing mass)
- Manley and Salesky performed an isobar-model partial-wave analysis of the world's available set of bubble-chamber data for these reactions many years ago [PRD **45**, 4002 (1992)]. Data set consisted of about 241,000 events (very low statistics by modern standards).
- Biggest problems began around 1600 MeV, where the number of important partial waves became greater than the data available to determine them. The amplitudes for quasi-two-body reactions as  $\pi N \rightarrow \pi \Delta$  and  $\pi N \rightarrow \rho N$  become quite noisy. A new isobar model analysis could be performed to determine the  $\pi \Delta$  and  $\rho N$  couplings of  $N^*$  and  $\Delta^*$  resonances more precisely.
- Such an analysis could incorporate the new data for  $\pi^- p \rightarrow \pi^0 \pi^0 n$ , which were measured at BNL by the Crystal Ball Collaboration.
- Needed to analyze data from Jefferson Lab in the reaction  $e p \rightarrow e' \pi^- \pi^+ p$ .
- An isobar model analysis is nontrivial, but can be done with new data.



## New baryons in the $\eta\Delta$ and $\omega\Delta$ channels

- The  $\eta\Delta$  channel is pure  $I=3/2$  and presents an opportunity to discover new  $\Delta^*$  resonances in these final states. The  $\eta\Delta$  channel could be studied by the reactions
  - $\pi^- p \rightarrow \pi^- \eta p$  (identify  $\eta$  by missing mass; select  $\Delta^0$  events by invariant mass of  $\pi^- p$ )
  - $\pi^+ p \rightarrow \pi^+ \eta p$  (identify  $\eta$  by missing mass; select  $\Delta^{++}$  events by invariant mass of  $\pi^+ p$ )
  - Isospin invariance of the strong interactions means that these two reactions must give consistent results for  $\eta\Delta$  couplings. This presents a tight constraint to make sure that the couplings are determined consistently.
- Similarly,  $\omega\Delta$  couplings are also pure  $I=3/2$  and could be studied by
  - $\pi^- p \rightarrow \pi^- \omega p$  (identify  $\omega$  by missing mass; select  $\Delta^0$  events by invariant mass of  $\pi^- p$ )
  - $\pi^+ p \rightarrow \pi^+ \omega p$  (identify  $\omega$  by missing mass; select  $\Delta^{++}$  events by invariant mass of  $\pi^+ p$ )
- These reactions present a good opportunity to search for new and missing  $\Delta^*$  resonances. Quark-model predictions have been made by Simon Capstick and Winston Roberts (PRD **57**, 4301 (1998)).

## Strangeness Production ( $\Sigma$ 's)

- $\pi^+ p \rightarrow K^+ \Sigma^+ \rightarrow K^+ \pi^+ n$ 
  - Detect neutron by missing mass, and reconstruct  $\Sigma^+$  by its invariant mass.
  - This reaction is especially important because it is pure  $I=3/2$  and will excite only  $\Delta^*$  resonances.
- $\pi^- p \rightarrow K^+ \Sigma^- \rightarrow K^+ \pi^- n$ 
  - Detect neutron by missing mass and reconstruct  $\Sigma^-$  by its invariant mass.
- $\pi^- p \rightarrow K^0 \Sigma^0$ 
  - $$\begin{array}{l} \downarrow \quad \downarrow \\ \quad \quad \downarrow \rightarrow \gamma \Lambda \rightarrow \gamma \pi^- p \\ \downarrow \rightarrow K_S \rightarrow \pi^+ \pi^- \end{array}$$
  - Detect  $\gamma$  by missing mass, reconstruct  $\Lambda$  from  $\pi p$  invariant mass, then reconstruct  $\Sigma$  from  $\gamma \Lambda$ .
- Previous two reactions are complementary, since they are a mixture of  $I=1/2$  and  $I=3/2$ . Both are needed.



# Strangeness Production

- Resonance couplings determined for  $K \Lambda$  and  $K \Sigma$  can be compared with quark model calculations, such as those of Simon Capstick and Winston Roberts (PRD **58**, 074011 (1998)).
- Quark-model predictions have also been made for other channels involving strange particles, such as  $K^* \Lambda$ ,  $K \Lambda (1405)$ , and  $K \Lambda (1520)$ .



## Recent Progress

New veto wall constructed

DAQ electronics upgraded

Magnet, TPC winding completed

Zip tracking hardware improved

## What is needed

Zip tracking of magnet

DAQ expertise

Electronics expertise

Hall probes, beam calibration

Install Plastic Ball as a back angle detector

Commissioning of new experiment

# Invitation

Additional collaborators are welcome!  
Contact me ([sadler@physics.acu.edu](mailto:sadler@physics.acu.edu))  
Or Rajendran Raja ([raja@fnal.gov](mailto:raja@fnal.gov))

Full MIPP proposal:

<http://ppd.fnal.gov/experiments/e907/notes/MIPPnotes/public/pdf/MIPP0138/MIPP0138.pdf>



# Summary of Present Status

- Listings in Review of Particle Properties rely on PWA's from 1980.
- GWU (formerly VPI) analyses fit data well, but other analyses using different approaches are also needed.
- Analyses should include  $\pi p \rightarrow K\Lambda$ ,  $\pi p \rightarrow \pi\pi N$  and other inelastic channels. New efforts by EBAC, Bonn, Zagreb and other groups are now in place. Effect on PDG?
- PWA for hadronic channels is needed to analyze data from photo- and electro-production of resonances.
- MIPP is a unique (existing!) experimental facility that can be used for baryon spectroscopy (1-3 GeV/c  $\pi$ 's).